

Congenital Arteriovenous Malformation Located in the Posterior Mediastinum

Arteriovenous Malformation

Ali Celik¹, İlknur Aytekin¹, Sevki Mustafa Demiröz¹, Özgür Ekinci², İsmail Cüneyt¹

¹Department of Thoracic Surgery,

²Department of Pathology, Gazi University Medical School, Ankara, Turkey

Abstract

Congenital arteriovenous malformations are not real tumor bodies and usually do not have endothelial proliferation. They arise from abnormal development of vascular structures and cause aberrant capillary network connections between arteries and veins. While the most usual sites of location for these pathologies are lower extremities and the neck, mediastinal location is very rare. A posterior mediastinum located lesion was found in the 50 years old patient examined due to back pain. Total excision performed and the pathologic examination revealed an arteriovenous malformation. In this article, we reviewed the clinical features and treatment approaches of the mediastinum located arteriovenous malformations in light of the literature.

Keywords

Arteriovenous Malformation; Posterior Mediastinum; Surgery

DOI:10.4328/ECAM.2

Received : 30.01.2013

Accepted : 07.03.2013

Published Online : 11.03.2013

Printed Online : 11.03.2013

Eu Clin Anal Med 2013;1(1): 18-20

Corresponding Author: Ali Celik, Gazi University School of Medicine, Department of Thoracic Surgery, Besevler, 06500, Ankara, Turkey.

GSM: +905304045467 • **F:** +903122025891 • **E-Mail:** alicelik78@gmail.com

How to cite this article: Ali Celik, İlknur Aytekin, Sevki Mustafa Demiroz, Muhammet Sayan, Ozgur Ekinci, İsmail Cüneyt Kurul. Congenital Arteriovenous Malformation Located in the Posterior Mediastinum. Eu Clin Anal Med 2013;1(1): 18-20

Introduction

Mediastinum located congenital arteriovenous malformations (AVM) are quite uncommon [1,4]. Although the AVMs generally seen in the childhood period are usually congenital, they may also be acquired [3,4]. The most feared complication of these pathologies occurring by the combination of the small arteriovenous fistula structures is bleeding. These lesions are usually asymptomatic and detected incidentally [1,2]. The lesions occupying space in the posterior mediastinum as well as the other pathologies resulting from the vascular structures should be considered in the differential diagnosis. Surgery is the treatment of choice considering the clinical features, complications and the differential diagnosis.

Case Report

A 50 years old male patient who was evaluated for the lower back pain, referred to our clinic for right hilar enlargement found on the pulmonary radiographs. The patient had no trauma, coughing and hemoptysis history and physical examination, vital symptoms, sedimentation rate and the other laboratory outcomes were in normal ranges. A soft tissue lesion of 23x43x43 mm in size causing cortical thinning and destruction in the right half of the T7 vertebra was found on the pulmonary computed tomography (CT) (Figure 1). There was no other pathology observed related to the lung parenchyma and mediastinal structures. On magnetic resonance (MRI) examination, lesion locating in the right paravertebral region was expanding to the paravertebral soft tissue plans, and had a heterogeneous signal featuring fat content. The lesion was smooth lobulated contoured, presenting a prominent infiltration to the adjacent pleura but was not prominently associated with the vascular structures. Patient underwent muscle sparing posterolateral thoracotomy. Lesion was observed in the exploration with a base in the vertebra corpus, destructing the corpus with a rich blood built up,




Figure 1. A soft tissue lesion of 23 x 43 x 43 mm with broken fat density that has an expansile character and could be compatible with hemangioma was observed in the right half of the T7 vertebra corpus.




Figure 2. Low power view of the vascular lesion with varying vessel wall thickness and luminal protrusions. H&E, x40 magnification (A). In a displaced aggregate, anastomosing vascular structures with different luminal calibers form an abnormal mesh. H&E, x40 magnification (B).

resembling a vessel glomus but has not a relation with the large vascular structures was excised, trimming the destructed bone tissue. The patient has not a problem in the postoperative period was discharged on the fourth day. The result of the pathology was reported as an arteriovenous malformation (Figure 2).

Discussion

AVMs are usually congenital, may also be acquired related to a penetrating trauma or iatrogenically developed lesions following various surgical interventions as needle biopsy, vascular catheterization or embolotomy [3,5]. These may be rarely seen in the chest wall or mediastinum. The blood from the arterial system usually passes to the venous system without distribution to the tissue because of the AVMs. Virtually the real pathology is arteriovenous fistula. Therefore, the tissue cannot be fed sufficiently, blood flow of the surrounding tissue is impaired and the changes resulting from the malnutrition are observed in the adjacent structures. This can be demonstrated through angiography in much larger AVMs, so called as steal phenomenon [2]. In fact, this physiopathological mechanism lies under all the events occurring due to AVMs. In our case also, the lesion led to destruction in the corpus of the adjacent vertebra.

The clinical feature of these mostly asymptomatic lesions depends on the size and location of the lesion. The pressure symptoms of the trachea, esophagus and heart as to the mediastinal location or the symptoms related to the bone involvement may be seen [4]. AVMs in the asymptomatic elderly patients those had not any characteristic in their history are usually congenital. Especially the AVMs in the neonatal and childhood period may be encountered with the symptoms and finding of the high-output heart failure [2]. Furthermore, the congenital AVMs may be accompanied by the other congenital diseases such as hereditary hemorrhagic telangiectasia (Osler-Weber Rendu). Pathological vessel connections in the mediastinal AVMs may be between the intercostal artery and vein, azygos vein or brachycephalic artery and vein and vena cava superior [1,2].

Primarily, CT is helpful for the diagnosis of the lesions. CT may suggest the exact location, nature and the relation with the surrounding structures of the lesion and the surgical technique will be performed. In the lesions preoperatively presumed as AVM, embolization is successfully applied with MRI angiography by the interventional radiologists. Particularly, the treatment of the larger lesions can be successfully done with the angiography and arterial embolization.

Besides the peripheral nerve sheath tumors commonly seen in this region, extrapulmonary sequestration and the mesenchymal tumors of vascular origin as hemangioma, angiolipoma, hemangioendothelioma and glomus tumors should be kept in mind for the differential diagnosis [2,6,7]. Hemangiomas, holding an important place in the differential diagnosis are accounted for nearly 90% of the benign vascular lesions are usually asymptomatic [6]. The hemangiomas may be found in each three compartment of the mediastinum, and they occasionally may be located in the mediastinum, leading to symptoms such as horner's syndrome and spinal cord compression. More contrastation is observed in the central of the tumor than the periphery on the CT [6]. Angiosarcomas and their variants are malignant lesions and usually may be originated from the main vascular structures such as the vena cava superior, brachiocephalic vein, aorta, heart and pericardium and these are aggressive coursed tumors [7]. Hemangioendotheliomas are vascular endotelial tumors originated from the medium-large venous structures (e.g. brachiocephalic vein, vena cava) in the mediastinum. Their clinical behavior may resemble the benign hemangiomas as well as malignant tumors, angiosarcomas. Angiolipomas are the benign tumors consisting of the blood vessels and mature adipose tissue those

usually located in the subcutaneous tissue and rarely in the mediastinum [7].

In these cases of the low surgical mortality and morbidity, complete excision of the lesions is possible. The most feared complications of the AVMs are the spontaneous rupture and hemorrhage [2]. Hemostasis and bleeding control is crucial in the patients for whom the surgery is planned. The surgery with the preoperative embolization can be done more safely in the cases considering the intraoperative bleeding control will be difficult [2,4]. Anti-angiogenic therapy, sclerotherapy and steroids or interferon treatment can be tried in the cases without complete surgery [8].

Scientific Responsibility Statement

The authors declare that they are responsible for the article's scientific content including study design, data collection, analysis and interpretation, writing, some of the main line, or all of the preparation and scientific review of the contents and approval of the final version of the article.

Animal and human rights statement

All procedures performed in this study were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. No animal or human studies were carried out by the authors for this article.

Funding: None

Conflict of interest

None of the authors received any type of financial support that could be considered potential conflict of interest regarding the manuscript or its submission.

References

1. Lunde P, Sörli D, Bolz KD, Eide TJ. Huge arteriovenous malformation in the mediastinum. Clinical presentation and surgical management. *Scand J Thorac Cardiovasc Surg.* 1984;18(1):75-80.
2. Tsitouridis I, Michaelides M, Pervana S. Posterior Mediastinal Arteriovenous Malformation. *J Thorac Imaging* 2008; 23(2):127-130
3. Creech O Jr, Gantt J, Wren H. Traumatic arteriovenous fistula at unusual sites. *Ann Surg.* 1965;161:908-20.
4. Tennyson C, Routledge T, Chambers A, Scarci M. Arteriovenous malformation in the anterior mediastinum. *Ann Thorac Surg* 2010; 90:e 9-10
5. Chloroyiannis Y, Reul JG. Iatrogenic Left Subclavian Artery-to-Left Brachiocephalic Vein Fistula Successful Repair without a Sternotomy. *Tex Heart Inst J* 2004;31:172-4
6. Hirai K, Takeuchi S, Bessho R, Ohaki Y, Koizumi K, Shimizu K. Venous hemangioma of the anterior mediastinum. *J Nippon Med Sch.* 2010;77(2):115-8.
7. Wada H, Teramatsu T. Mediastinal tumors-a statistical nationwide report of 1,546 cases between July, 1975 and May, 1979 in Japan. *Nippon Kyobu Geka Gakkai Zasshi* 1982;30:374-378
8. Fishman SJ. Vascular anomalies of the mediastinum. *Semin Pediatr Surg.* 1999;8:92-8.